



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2009

ER , ER , and gpER: novel aspects of oestrogen receptor signalling in atherosclerosis

Meyer, M R ; Barton, M

DOI: <https://doi.org/10.1093/cvr/cvp187>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-32167>

Journal Article

Published Version

Originally published at:

Meyer, M R; Barton, M (2009). ER , ER , and gpER: novel aspects of oestrogen receptor signalling in atherosclerosis. *Cardiovascular Research*, 83(4):605-610.

DOI: <https://doi.org/10.1093/cvr/cvp187>

ER α , ER β , and gpER: novel aspects of oestrogen receptor signalling in atherosclerosis

Matthias R. Meyer and Matthias Barton*

Departement für Innere Medizin, Klinik und Poliklinik für Innere Medizin, Universitätsspital Zürich, Rämistrasse 100, CH-8091 Zürich, Switzerland

Online publish-ahead-of-print 18 June 2009

Cardiovascular disease represents the leading cause of morbidity and mortality in women. Epidemiological and experimental evidence indicates several atheroprotective effects of endogenous oestrogens, which interfere with atherosclerosis progression and inflammation. Nevertheless, oestrogen receptor function across different stages of atherosclerosis development remains poorly characterized. This viewpoint editorial discusses the current knowledge of oestrogen receptor function in physiology and within different stages of atherogenesis. Increasing our understanding of the molecular mechanisms determining oestrogen action in humans, with and without established atherosclerosis, may help to develop new strategies for the treatment of cardiovascular disease in women, and possibly also in men.

1. Endogenous oestrogens and cardiovascular risk: role of menopause

Despite substantial efforts to improve education and public awareness and despite the use of effective medications and life-style changes for controlling the associated risk factors—hypertension, tobacco use, hypercholesterolemia, obesity, and diabetes—cardiovascular disease remains the leading cause of death in women worldwide.^{1,2} In contrast to age-matched men, the incidence of clinical manifestations of coronary artery disease is considerably lower in premenopausal women; indeed, about 95% of women develop cardiovascular disease after menopause when endogenous oestrogen levels are low.^{3–5} Accordingly, endogenous oestrogen protects from accelerated atherogenesis due to obesity in female teenagers and young adults compared with their male counterparts.⁶ Cardiovascular risk increases after bilateral ovariectomy and in conditions associated with impaired ovarian function.^{3,7} Thus, ovarian dysfunction and either natural or surgical menopause have been recognized as a major risk factor for accelerated atherosclerotic vascular disease development. In addition, cardiovascular risk factors and certain polymorphisms in oestrogen receptor genes may also determine an earlier onset of menopause.⁸

The opinions expressed in this article are not necessarily those of the Editors of *Cardiovascular Research* or of the European Society of Cardiology.

* Corresponding author. Tel: +41 44 255 56 63; fax: +41 44 255 87 47.
E-mail address: matthiasbarton@gmx.com

Given that the number of postmenopausal women is expected to rise by 1 billion worldwide within the next 40 years,⁹ understanding the impact of menopause on women's cardiovascular health has become increasingly important.

In addition to the epidemiological evidence, numerous *in vitro* studies have identified mechanisms whereby oestrogens exert beneficial effects on the cardiovascular system.^{10,11} *In vivo*, the presence of endogenous oestrogens and their effect on cardiovascular homeostasis appear to be closely related to the degree of atherosclerosis progression throughout a woman's life.¹² Reversible fatty streak lesions are present *in utero*,¹³ and 13% of US women aged 30–34 years already display advanced coronary artery lesions susceptible to rupture.¹⁴ In stages of disrupted ovulatory cycling, low levels of endogenous oestrogens during premenopausal years accelerate the progression of atherosclerosis,^{12,15} which can be reversed by oestrogen therapy in animals.¹² In addition, results from experimental studies and recent clinical trials indicate that oestrogen therapy started within few years after menopause, i.e. before the development of severe atherosclerosis, may in fact reduce cardiovascular risk.^{5,16–20} In contrast, initiation of oestrogen therapy many years after menopause, i.e. when advanced and multiple atherosclerotic lesions are present, may have no or even deleterious cardiovascular effects.^{5,16–20} Thus, the vascular response to endogenous and exogenous oestrogens appears to change with ageing and the presence or progression of atherosclerosis: while oestrogen could prevent the development of atherosclerotic plaques in newly menopausal women, it would have no or even harmful effects in older postmenopausal women with advanced atherosclerotic disease and vulnerable plaques. This has been referred to as the 'timing hypothesis'.^{5,16–20}

2. Vascular oestrogen receptor signalling in women and men

Biological effects of oestrogens are mediated by three oestrogen receptors: oestrogen receptor- α (ER α), oestrogen receptor- β (ER β), and the intracellular, transmembrane G protein-coupled oestrogen receptor (gpER).²¹ The 'classical' ER α and ER β receptors act as ligand-activated transcription factors that reside in the cytosol and translocate into the nucleus upon ligand binding. Subsequently, oestrogen

receptors interact with oestrogen response elements in the promoter region of target genes.¹¹ In addition, oestrogens bind to plasma membrane-associated subpopulations of ER α and ER β , thereby activating a variety of rapid intracellular signalling cascades.^{21,22}

Oestrogens also activate gpER, previously termed GPR30, which is located to the endoplasmic reticulum and mediates rapid oestrogen signalling.²³ gpER is widely expressed in human tissues, including the cardiovascular system.^{24,25} Interestingly, oestrogen signalling in cardiomyocytes involves ER α - and ER β -independent pathways,²⁶ and treatment with the oestrogen receptor antagonists ICI 182,780 and tamoxifen, which are agonists of gpER,²⁷ inhibits cardiac cell growth.²⁸ Although genetic linkage studies indicated a potential role of the gpER locus on human chromosome 7p22 in the susceptibility to low-renin hypertension,²⁹ the role of gpER for vascular homeostasis remained unclear. We recently found that selective gpER activation acutely dilates human arteries, and markedly reduces blood pressure even under normotensive conditions, indicating that this receptor indeed controls vascular tone.³⁰ Moreover, selective gpER activation potentially inhibits human vascular smooth muscle cell growth,³⁰ (Figure 1) consistent with the growth-inhibitory effect of ICI 182,780 and tamoxifen in cardiomyocytes.²⁸ The clinical significance of gpER as a mediator of the vascular protective effects of oestrogens in humans remains to be determined;²¹ however, a recent subanalysis of the Raloxifene Use for The Heart (RUTH) trial suggests that treatment with a gpER agonist (raloxifene) may reduce cardiovascular risk in younger postmenopausal women.³¹

ER α , ER β , and gpER are expressed in the arterial wall of both women and men,^{11,25} and the non-selective oestrogen receptor agonist 17 β -estradiol has potent dilator effects on vascular tone of human coronary and internal mammary arteries obtained from either female or male patients.^{25,32}




Figure 1 Involvement of gpER in regulation of vascular function, vascular smooth muscle cell growth, and blood pressure. The non-selective oestrogen receptor agonist 17 β -estradiol (E2) and the selective gpER agonist G-1 acutely induce a relaxant response in human internal mammary arteries when compared with solvent control (CTL). However, the dilator effect of G-1 is even more potent than E2 (A). gpER activation also potentially and concentration-dependently inhibits serum-stimulated cell proliferation in human vascular smooth muscle cells (B). Intravenous infusion of G-1 at increasing concentrations reduces mean arterial blood pressure (MAP) in normotensive Sprague-Dawley rats. For comparison, the pressure response to acetylcholine (ACh) is shown (C). Figure reproduced in part from Haas, E., Bhattacharya, I., Brailoiu, E., Damjanovic, M., Brailoiu, G.C., Gao, X., Mueller-Guerre, L., Marjon, N.A., Gut, A., Minotti, R., Meyer, M.R., Amann, K., Ammann, E., Perez-Dominguez, A., Genoni, M., Clegg, D.J., Dun, N.J., Resta, T.C., Prossnitz, E.R., Barton, M., 2009. Regulatory role of G protein-coupled estrogen receptor for vascular function and obesity. *Circulation Research*. 104, 288–291. © 2009 American Heart Association.

These findings suggest a potential role for oestrogen receptor function also in the male cardiovascular system. Interestingly, 80% of plasma-bound 17 β -estradiol in men originates from aromatization of testosterone and androstenedione in peripheral tissues by the enzyme aromatase,³³ which is highly expressed in the male vasculature.³⁴ Moreover, aromatase is expressed at high levels in the endoplasmic reticulum,³⁵ to which gpER is located.²³ Pharmacological inhibition of aromatase accelerates the formation of atherosclerotic lesions in male animals,³⁶ and impairs flow-mediated vasodilation in young men.³⁷ Moreover, male mice lacking functional ER β develop hypertension.³⁸ In line with the finding that ER α mediates atheroprotective effects,³⁹ impaired vascular function and premature coronary artery disease were noted in a man with a disruptive mutation in the ER α gene.⁴⁰ Thus, not only the female but also the male cardiovascular system appears to be an important source and target for oestrogens affecting vascular disease development.^{11,41} Nevertheless, studies in humans comparing oestrogen plasma concentrations and the progression of cardiovascular disease revealed conflicting results.⁴¹ At present, there is doubt that treatment of male patients with oestrogen receptor-activating compounds, as shown experimentally in mice⁴² and previously unsuccessfully attempted in male patients,^{43,44} represents a treatment option to interfere with atherosclerosis progression. However, 17 β -estradiol has been successfully used in male patients with coronary artery disease as a coating agent for endovascular stents, as indicated by the promising results of the Estrogen And Stents To Eliminate Restenosis (EASTER) trial.⁴⁵

3. Does the development of atherosclerosis affect oestrogen receptor function?

Experimental studies suggest that at least in mice ER α appears to be largely responsible for the protective effects of oestrogens against atherosclerotic vascular disease.³⁹ On the other hand, expression of ER β in humans correlates with coronary calcification and atherosclerosis independent of age.⁴⁶ Indeed, ER β is expressed at higher levels than ER α in postmenopausal women with advanced atherosclerosis.⁴⁶ Moreover, ER β expression is up-regulated following balloon angioplasty of structurally normal arteries of mice.⁴⁷ However, the absence of atherosclerosis makes the clinical relevance of this experimental model questionable. Interestingly, ER β exhibits an inhibitory action on ER α -dependent gene expression and may oppose the actions of ER α , including its effects on facilitating nitric oxide-dependent, endothelium-mediated relaxation.^{48,49} This could explain the loss of oestrogen responsiveness of arteries with atherosclerosis. Conversely, both ER β deficiency in laboratory animals as well as functionally relevant mutations of the ER β gene in women have been linked to the development of hypertension,^{38,50} and associations of two ER β polymorphisms with an increased cardiovascular risk in women have been reported.⁵¹ It has also been recently demonstrated that ER α and ER β regulate distinct and largely non-overlapping sets of genes in the arterial vascular wall.⁵² Thus, both ER α and ER β may have distinct roles ensuring vascular homeostasis that may, however, be affected by the presence of atherosclerosis.

According to some studies, the abundance of both oestrogen receptor subtypes ER α and ER β in the aorta of humans decreases with the progression of atherosclerosis.^{53,54} Inactivation of ER α and ER β genes in the vasculature may be caused by DNA methylation of the promoter region of these genes; DNA methylation, an important epigenetic mechanism, increases with ageing and plaque progression.^{55–57} Thus, structural changes of oestrogen receptors may result in functional loss of responsiveness to oestrogens. Indeed, proliferating vascular smooth muscle cells, which have undergone phenotypic modulation upon migrating to the intima,⁵⁸ display a high degree of ER α methylation.⁵⁶ However, we currently do not know whether and by which mechanisms changes in oestrogen receptor expression and structure in vascular cells affect atherogenesis, or whether vascular cells without functional oestrogen receptors become more abundant and prevent effects of oestrogens from becoming functional.

It has been suggested that the local formation of biologically active estrogens is higher in human aortas with mild atherosclerotic changes than in those with more advanced lesions.⁵⁹ Moreover, the ER β -associated co-regulatory protein NM23-H2, which activates ER β signalling *in vitro*, is down-regulated with progression of atherosclerosis in coronary arteries.⁶⁰ The notion that the protective effects of vascular oestrogen signalling are lost with the progression of atherosclerosis is further supported by the finding that 27-hydroxycholesterol, an abundant cholesterol metabolite found in atherosclerotic lesions, acts as an oestrogen receptor antagonist.⁶¹ In addition, the aforementioned inactivation of oestrogen receptor genes, an altered balance of ER α and ER β protein expression, and changes in oestrogen receptor signalling during different steps of atherogenesis

may add up to a different regulation of oestrogen-sensitive genes, as well as of vascular homeostasis in general (Figure 2). Oestrogen therapy may also have beneficial effects on the expression and function of genes regulating calcium homeostasis that are limited to early, but not advanced, stages of atherosclerotic lesions.¹⁹ Moreover, oestrogen-dependent induction of matrix metalloproteinases, proteolytic enzymes capable of degrading components of the extracellular matrix expressed in fibrous caps, may contribute to maintenance of lumen size in early atherosclerosis, but could increase the likelihood of plaque erosion or rupture in more advanced stages of the disease.^{19,62}

Animal studies by Oparil's group have found that—unlike in young ovariectomized rats—oestrogen therapy in aged animals fails to prevent neointima formation and inflammatory responses following balloon angioplasty in structurally normal arteries.⁶³ This indicates that ageing—even in the absence of atherosclerosis—appears to modulate the chronic vascular response to natural oestrogens.¹⁸ It has also been suggested that oestrogens, under certain conditions such as prolonged oestrogen deficiency, and particularly when used in the context of vascular inflammation as seen in atherosclerosis, may attenuate vasodilator capacity, enhance inflammatory activity, and improve plaque instability.^{64,65} Accordingly, a prolonged period of hypoestrogenicity disrupts both neuroprotective and anti-inflammatory actions of oestrogens in animals.⁶⁶ In addition, oestrogen-dependent direct and endothelium-mediated relaxation as well as up-regulation of ER α and eNOS protein levels in ovariectomized rats is maintained when 17 β -estradiol treatment is initiated 4 months, but not 8 months after surgery.⁶⁷ Together, these experimental studies provide evidence to support the concept of the 'timing hypothesis'.




Figure 2 Molecular targets of oestrogens in the normal (left) and atherosclerotic vascular wall (right). With the progression of an atherosclerotic lesion as well as with ageing, expression of ER α and ER β becomes a target of methylation (CH₃)-associated inactivation. Moreover, expression of oestrogen receptor-associated co-regulatory proteins, such as NM23-H2, is altered, and ER β expression changes in advanced atherosclerosis. Local formation of biologically active oestrogens through aromatase and other pathways appears to be reduced (E₂ ↓) under these conditions. Based on these changes of oestrogen receptor signalling in advanced atherosclerosis, it is currently not clear how endogenous oestrogens affect oestrogen-dependent proteins involved in inflammation and structure of the atherosclerotic plaque, such as inflammatory nitric oxide synthase (iNOS) and matrix metalloproteinases (MMP). This illustrates the complexity of action of oestrogens and how equine oestrogen mixtures containing hormonal substances of unidentified activity affect oestrogen receptor function and regulation of oestrogen-sensitive genes in the presence of atherosclerosis. gpER, G protein-coupled estrogen receptor.

4. Clinical implications and perspectives

Although there is now substantial evidence suggesting that vascular functionality of oestrogen receptors depends on the length of a period of hypoestrogenicity and the degree of atherosclerosis progression, it is still not known how the cessation of endogenous oestrogen production after menopause affects oestrogen receptor signalling in human vascular cells *in vivo*. Therefore, future basic research should include studies that will help identifying the nature of changes of oestrogen receptor function in diseased arteries. Such changes were not considered in earlier randomized hormone therapy trials such as the Heart and Estrogen/progestin Replacement Study (HERS) and the Women's Health Initiative (WHI), which demonstrated adverse cardiovascular outcomes of hormone therapy in women late after menopause.^{68,69} The medical regimen used in these studies consisted of conjugated equine oestrogens derived from horse urine, and of synthetic progestins such as medroxyprogesterone acetate.^{68,69} Unlike 17 β -estradiol, the major endogenous human oestrogen lost after menopause, conjugated equine estrogens contain a mixture of at least 10 different oestrogens, several androgens, progestins, and other substances of yet unknown vascular and procoagulatory activity.^{5,18,41} These substances display different or even unknown selectivity and binding affinity for oestrogen receptors,⁵ which result in unpredictable effects on multiple oestrogen signalling pathways evoking random vascular responses *in vivo*, especially in the presence of atherosclerosis. Moreover, medroxyprogesterone acetate has been reported to abrogate beneficial effects of oestrogens on vascular smooth muscle cell growth, lipid profile, and vascular function, even turning oestrogen-induced vasodilation into coronary vasospasm.^{70–72}

Currently, prospective clinical trials are underway to test whether natural oestrogen has therapeutic potential in women with atherosclerosis: The Kronos Early Estrogen Protection Study (KEEPS; ClinicalTrials.gov number, NCT00154180), and the Early vs. Late Intervention Trial With Estradiol (ELITE; ClinicalTrials.gov number, NCT00114517). Moreover, the use of natural oestrogens such as 17 β -estradiol at low doses, and possibly intermittent administration mimicking the normal menstrual cycle,^{18,41} potentially will help to preserve activating signalling cascades, which are responsible for the beneficial vascular effects of oestrogens in premenopausal women. Finally, the use of selective oestrogen receptor modulators such as raloxifene, which act as gpER agonists,²⁷ might be suitable for treatment and prevention of atherosclerotic vascular disease. Results from the RUTH trial support this notion, at least in younger postmenopausal women.³¹

At present, hormone therapy is not an option for primary or secondary prevention of cardiovascular disease in women. There are proven therapies such as simple changes in lifestyle including maintaining a normal body weight, a diet rich in fruits and vegetables, and regular physical exercise. These measures—although still underused—should be initiated as early as possible to reduce the burden of atherosclerotic vascular disease later in life.^{2,18}

Conflict of interest: none declared.

Funding

Original work by the authors is supported by the Swiss National Science Foundation [grant numbers 3200-108258/1, and K-33KO_122504/1].

References

1. Yusuf S, Reddy S, Ounpuu S, Anand S. Global burden of cardiovascular diseases: part I: general considerations, the epidemiologic transition, risk factors, and impact of urbanization. *Circulation* 2001;**104**:2746–2753.
2. Libby P. The forgotten majority: unfinished business in cardiovascular risk reduction. *J Am Coll Cardiol* 2005;**46**:1225–1228.
3. Witteman JC, Grobbee DE, Kok FJ, Hofman A, Valkenburg HA. Increased risk of atherosclerosis in women after the menopause. *Br Med J* 1989;**298**:642–644.
4. Barrett-Connor E. Sex differences in coronary heart disease. Why are women so superior? The 1995 Ancel Keys Lecture. *Circulation* 1997;**95**:252–264.
5. Dubey RK, Imthurn B, Barton M, Jackson EK. Vascular consequences of menopause and hormone therapy: importance of timing of treatment and type of estrogen. *Cardiovasc Res* 2005;**66**:295–306.
6. McGill HC Jr, McMahan CA, Herderick EE, Zieske AW, Malcom GT, Tracy RE *et al*. Obesity accelerates the progression of coronary atherosclerosis in young men. *Circulation* 2002;**105**:2712–2718.
7. Punnonen R, Jokela H, Aine R, Teisala K, Salomaki A, Uppa H. Impaired ovarian function and risk factors for atherosclerosis in premenopausal women. *Maturitas* 1997;**27**:231–238.
8. Bittner V. Menopause and cardiovascular risk cause or consequence? *J Am Coll Cardiol* 2006;**47**:1984–1986.
9. Barton M, Meyer MR. Postmenopausal hypertension: mechanisms and therapy. *Hypertension* 2009;**54**:11–18.
10. Mendelsohn ME, Karas RH. The protective effects of estrogen on the cardiovascular system. *N Engl J Med* 1999;**340**:1801–1811.
11. Meyer MR, Haas E, Barton M. Gender differences of cardiovascular disease: new perspectives for estrogen receptor signaling. *Hypertension* 2006;**47**:1019–1026.
12. Clarkson TB. Estrogen effects on arteries vary with stage of reproductive life and extent of subclinical atherosclerosis progression. *Menopause* 2007;**14**:373–384.
13. Napoli C, D'Armiento FP, Mancini FP, Postiglione A, Witztum JL, Palumbo G *et al*. Fatty streak formation occurs in human fetal aortas and is greatly enhanced by maternal hypercholesterolemia. Intimal accumulation of low density lipoprotein and its oxidation precede monocyte recruitment into early atherosclerotic lesions. *J Clin Invest* 1997;**100**:2680–2690.
14. McMahan CA, Gidding SS, Fayad ZA, Zieske AW, Malcom GT, Tracy RE *et al*. Risk scores predict atherosclerotic lesions in young people. *Arch Intern Med* 2005;**165**:883–890.
15. Baird Merz CN, Johnson BD, Sharaf BL, Bittner V, Berga SL, Braunstein GD *et al*. Hypoestrogenemia of hypothalamic origin and coronary artery disease in premenopausal women: a report from the NHLBI-sponsored WISE study. *J Am Coll Cardiol* 2003;**41**:413–419.
16. Salpeter SR, Walsh JM, Greyber E, Salpeter EE. Brief report: coronary heart disease events associated with hormone therapy in younger and older women. A meta-analysis. *J Gen Intern Med* 2006;**21**:363–366.
17. Rossouw JE, Prentice RL, Manson JE, Wu L, Barad D, Barnabei VM *et al*. Postmenopausal hormone therapy and risk of cardiovascular disease by age and years since menopause. *J Am Med Assoc* 2007;**297**:1465–1477.
18. Barton M, Meyer MR, Haas E. Hormone replacement therapy and atherosclerosis in postmenopausal women: does aging limit therapeutic benefits? *Arterioscler Thromb Vasc Biol* 2007;**27**:1669–1672.
19. Mendelsohn ME, Karas RH. HRT and the young at heart. *N Engl J Med* 2007;**356**:2639–2641.
20. Manson JE, Bassuk SS. Invited commentary: hormone therapy and risk of coronary heart disease why renew the focus on the early years of menopause? *Am J Epidemiol* 2007;**166**:511–517.
21. Meyer MR, Haas E, Prossnitz ER, Barton M. Non-genomic regulation of vascular cell function and growth by estrogen. *Mol Cell Endocrinol* 2009;**308**:9–16.
22. Moriarty K, Kim KH, Bender JR. Minireview: estrogen receptor-mediated rapid signaling. *Endocrinology* 2006;**147**:5557–5563.
23. Revankar CM, Cimino DF, Sklar LA, Arterburn JB, Prossnitz ER. A transmembrane intracellular estrogen receptor mediates rapid cell signaling. *Science* 2005;**307**:1625–1630.

24. Owman C, Blay P, Nilsson C, Lolait SJ. Cloning of human cDNA encoding a novel heptahelix receptor expressed in Burkitt's lymphoma and widely distributed in brain and peripheral tissues. *Biochem Biophys Res Commun* 1996;**228**:285–292.
25. Haas E, Meyer MR, Schurr U, Bhattacharya I, Minotti R, Nguyen HH *et al.* Differential effects of 17 β -estradiol on function and expression of estrogen receptor alpha, estrogen receptor beta, and GPR30 in arteries and veins of patients with atherosclerosis. *Hypertension* 2007;**49**:1358–1363.
26. Ullrich ND, Krust A, Collins P, MacLeod KT. Genomic deletion of estrogen receptors ERalpha and ERbeta does not alter estrogen-mediated inhibition of Ca²⁺ influx and contraction in murine cardiomyocytes. *Am J Physiol Heart Circ Physiol* 2008;**294**:H2421–H2427.
27. Thomas P, Pang Y, Filardo EJ, Dong J. Identity of an estrogen membrane receptor coupled to a G protein in human breast cancer cells. *Endocrinology* 2005;**146**:624–632.
28. Mercier I, Mader S, Calderone A. Tamoxifen and ICI 182,780 negatively influenced cardiac cell growth via an estrogen receptor-independent mechanism. *Cardiovasc Res* 2003;**59**:883–892.
29. Lafferty AR, Torpy DJ, Stowasser M, Taymans SE, Lin JP, Huggard P *et al.* A novel genetic locus for low renin hypertension: familial hyperaldosteronism type II maps to chromosome 7 (7p22). *J Med Genet* 2000;**37**:831–835.
30. Haas E, Bhattacharya I, Brailoiu E, Damjanovic M, Brailoiu GC, Gao X *et al.* Regulatory role of G protein-coupled estrogen receptor for vascular function and obesity. *Circ Res* 2009;**104**:288–291.
31. Collins P, Mosca L, Geiger MJ, Grady D, Kornitzer M, Amewou-Atiso MG *et al.* Effects of the selective estrogen receptor modulator raloxifene on coronary outcomes in the Raloxifene Use for The Heart trial: results of subgroup analyses by age and other factors. *Circulation* 2009;**119**:922–930.
32. Mügge A, Riedel M, Barton M, Kuhn M, Lichtlen PR. Endothelium independent relaxation of human coronary arteries by 17 beta-oestradiol in vitro. *Cardiovasc Res* 1993;**27**:1939–1942.
33. Liu PP, Fukuoka M. Sex hormones as novel risk biomarkers for atherosclerosis in peripheral vascular disease. *J Am Coll Cardiol* 2007;**50**:1077–1079.
34. Murakami H, Harada N, Sasano H. Aromatase in atherosclerotic lesions of human aorta. *J Steroid Biochem Mol Biol* 2001;**79**:67–74.
35. Hojo Y, Hattori TA, Enami T, Furukawa A, Suzuki K, Ishii HT *et al.* Adult male rat hippocampus synthesizes estradiol from pregnenolone by cytochromes P45017alpha and P450 aromatase localized in neurons. *Proc Natl Acad Sci USA* 2004;**101**:865–870.
36. Nathan L, Shi W, Dinh H, Mukherjee TK, Wang X, Lusis AJ *et al.* Testosterone inhibits early atherogenesis by conversion to estradiol: critical role of aromatase. *Proc Natl Acad Sci USA* 2001;**98**:3589–3593.
37. Lew R, Komesaroff P, Williams M, Dawood T, Sudhir K. Endogenous estrogens influence endothelial function in young men. *Circ Res* 2003;**93**:1127–1133.
38. Zhu Y, Bian Z, Lu P, Karas RH, Bao L, Cox D *et al.* Abnormal vascular function and hypertension in mice deficient in estrogen receptor beta. *Science* 2002;**295**:505–508.
39. Hodgins JB, Kregge JH, Reddick RL, Korach KS, Smithies O, Maeda N. Estrogen receptor alpha is a major mediator of 17 β -estradiol's atheroprotective effects on lesion size in ApoE^{−/−} mice. *J Clin Invest* 2001;**107**:333–340.
40. Sudhir K, Chou TM, Chatterjee K, Smith EP, Williams TC, Kane JP *et al.* Premature coronary artery disease associated with a disruptive mutation in the estrogen receptor gene in a man. *Circulation* 1997;**96**:3774–3777.
41. Meyer MR, Haas E, Barton M. Need for research on estrogen receptor function: importance for postmenopausal hormone therapy and atherosclerosis. *Gen Med* 2008;**5**(Suppl. A):S19–S33.
42. Bourassa PA, Milos PM, Gaynor BJ, Breslow JL, Aiello RJ. Estrogen reduces atherosclerotic lesion development in apolipoprotein E-deficient mice. *Proc Natl Acad Sci USA* 1996;**93**:10022–10027.
43. Marmorston J, Magdison O, Kuzma O, Moore FJ. Estrogen therapy in men with myocardial infarction. Side-effects with increasing dosage and time. *J Am Med Assoc* 1960;**174**:241–244.
44. Stamler J, Pick R, Katz LN, Pick A, Kaplan BM, Berkson DM *et al.* Effectiveness of estrogens for therapy of myocardial infarction in middle-age men. *J Am Med Assoc* 1963;**183**:632–638.
45. Abizaid A, Albaladejo M, Costa MA, Abizaid AS, Staico R, Feres F *et al.* First human experience with the 17-beta-estradiol-eluting stent: the Estrogen And Stents To Eliminate Restenosis (EASTER) trial. *J Am Coll Cardiol* 2004;**43**:1118–1121.
46. Christian RC, Liu PY, Harrington S, Ruan M, Miller VM, Fitzpatrick LA. Intimal estrogen receptor (ER)beta, but not ERalpha expression, is correlated with coronary calcification and atherosclerosis in pre- and postmenopausal women. *J Clin Endocrinol Metab* 2006;**91**:2713–2720.
47. Lindner V, Kim SK, Karas RH, Kuiper GG, Gustafsson JA, Mendelsohn ME. Increased expression of estrogen receptor-beta mRNA in male blood vessels after vascular injury. *Circ Res* 1998;**83**:224–229.
48. Matthews J, Gustafsson JA. Estrogen signaling: a subtle balance between ER alpha and ER beta. *Mol Interv* 2003;**3**:281–292.
49. Traupe T, Stettler CD, Li H, Haas E, Bhattacharya I, Minotti R *et al.* Distinct roles of estrogen receptors alpha and beta mediating acute vasodilation of epicardial coronary arteries. *Hypertension* 2007;**49**:1364–1370.
50. Ogawa S, Emi M, Shiraki M, Hosoi T, Ouchi Y, Inoue S. Association of estrogen receptor beta (ESR2) gene polymorphism with blood pressure. *J Hum Genet* 2000;**45**:327–330.
51. Rexrode KM, Ridker PM, Hegener HH, Buring JE, Manson JE, Zee RY. Polymorphisms and haplotypes of the estrogen receptor-beta gene (ESR2) and cardiovascular disease in men and women. *Clin Chem* 2007;**53**:1749–1756.
52. O'Lone R, Knorr K, Jaffe IZ, Schaffer ME, Martini PG, Karas RH *et al.* Estrogen receptors alpha and beta mediate distinct pathways of vascular gene expression, including genes involved in mitochondrial electron transport and generation of reactive oxygen species. *Mol Endocrinol* 2007;**21**:1281–1296.
53. Losordo DW, Kearney M, Kim EA, Jekanowski J, Isner JM. Variable expression of the estrogen receptor in normal and atherosclerotic coronary arteries of premenopausal women. *Circulation* 1994;**89**:1501–1510.
54. Nakamura Y, Suzuki T, Miki Y, Tazawa C, Senzaki K, Moriya T *et al.* Estrogen receptors in atherosclerotic human aorta: inhibition of human vascular smooth muscle cell proliferation by estrogens. *Mol Cell Endocrinol* 2004;**219**:17–26.
55. Post WS, Goldschmidt-Clermont PJ, Wilhide CC, Heldman AW, Sussman MS, Ouyang P *et al.* Methylation of the estrogen receptor gene is associated with aging and atherosclerosis in the cardiovascular system. *Cardiovasc Res* 1999;**43**:985–991.
56. Ying AK, Hassanain HH, Roos CM, Smiraglia DJ, Issa JJ, Michler RE *et al.* Methylation of the estrogen receptor-alpha gene promoter is selectively increased in proliferating human aortic smooth muscle cells. *Cardiovasc Res* 2000;**46**:172–179.
57. Kim J, Kim JY, Song KS, Lee YH, Seo JS, Jelinek J *et al.* Epigenetic changes in estrogen receptor beta gene in atherosclerotic cardiovascular tissues and in-vitro vascular senescence. *Biochim Biophys Acta* 2007;**1772**:72–80.
58. Campbell GR, Campbell JH, Manderson JA, Horrigan S, Rennick RE. Arterial smooth muscle. A multifunctional mesenchymal cell. *Arch Pathol Lab Med* 1988;**112**:977–986.
59. Nakamura Y, Suzuki T, Sasano H. Estrogen actions and in situ synthesis in human vascular smooth muscle cells and their correlation with atherosclerosis. *J Steroid Biochem Mol Biol* 2005;**93**:263–268.
60. Rayner K, Chen YX, Hibbert B, White D, Miller H, Postel EH *et al.* NM23-H2, an estrogen receptor beta-associated protein, shows diminished expression with progression of atherosclerosis. *Am J Physiol Regul Integr Comp Physiol* 2007;**292**:R743–R750.
61. Umetani M, Domoto H, Gormley AK, Yuhanna IS, Cummins CL, Javitt NB *et al.* 27-Hydroxycholesterol is an endogenous SERM that inhibits the cardiovascular effects of estrogen. *Nat Med* 2007;**13**:1185–1192.
62. Libby P, Aikawa M. Stabilization of atherosclerotic plaques: new mechanisms and clinical targets. *Nat Med* 2002;**8**:1257–1262.
63. Miller AP, Xing D, Feng W, Fintel M, Chen YF, Oparil S. Aged rats lose vasoprotective and anti-inflammatory actions of estrogen in injured arteries. *Menopause* 2007;**14**:251–260.
64. Mendelsohn ME, Karas RH. Molecular and cellular basis of cardiovascular gender differences. *Science* 2005;**308**:1583–1587.
65. Cann JA, Register TC, Adams MR, St Clair RW, Espeland MA, Williams JK. Timing of estrogen replacement influences atherosclerosis progression and plaque leukocyte populations in ApoE^{−/−} mice. *Atherosclerosis* 2008 doi:10.1016/j.atherosclerosis.2008.01.018.
66. Suzuki S, Brown CM, Dela Cruz CD, Yang E, Bridwell DA, Wise PM. Timing of estrogen therapy after ovariectomy dictates the efficacy of its neuroprotective and antiinflammatory actions. *Proc Natl Acad Sci USA* 2007;**104**:6013–6018.
67. Pinna C, Cignarella A, Sanvito P, Pelosi V, Bolego C. Prolonged ovarian hormone deprivation impairs the protective vascular actions of estrogen receptor alpha agonists. *Hypertension* 2008;**51**:1210–1217.

68. Hulley S, Grady D, Bush T, Furberg C, Herrington D, Riggs B *et al.* Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women. Heart and Estrogen/progestin Replacement Study (HERS) Research Group. *J Am Med Assoc* 1998;**280**: 605–613.
69. Rossouw JE, Anderson GL, Prentice RL, LaCroix AZ, Kooperberg C, Stefanick ML *et al.* Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results From the Women's Health Initiative randomized controlled trial. *J Am Med Assoc* 2002; **288**:321–333.
70. Effects of estrogen or estrogen/progestin regimens on heart disease risk factors in postmenopausal women. The Postmenopausal Estrogen/Progestin Interventions (PEPI) Trial. The Writing Group for the PEPI Trial. *J Am Med Assoc* 1995;**273**:199–208.
71. Miyagawa K, Rosch J, Stanczyk F, Hermsmeyer K. Medroxyprogesterone interferes with ovarian steroid protection against coronary vasospasm. *Nat Med* 1997;**3**:324–327.
72. Dubey RK, Jackson EK, Gillespie DG, Zacharia LC, Wunder D, Imthurn B *et al.* Medroxyprogesterone abrogates the inhibitory effects of estradiol on vascular smooth muscle cells by preventing estradiol metabolism. *Hypertension* 2008;**51**:1197–1202.